

Übungen zu Numerische Methoden II

Exercise Sheet 2

1. Consider the symmetric matrix

$$A = \begin{bmatrix} 4 & 1 & 0 & 0 & 0 \\ 1 & 4 & 1 & 0 & 0 \\ 0 & 1 & 4 & 1 & 0 \\ 0 & 0 & 1 & 4 & 1 \\ 0 & 0 & 0 & 1 & 4 \end{bmatrix}.$$

Localize the spectrum of the inverse matrix A^{-1} , using the Gershgorin circle theorem.

2. A matrix $A \in \mathbb{R}^{n \times n}$ is diagonalizable if it has n distinct eigenvalues. Apply the Gershgorin circle theorem to show that the following matrix is diagonalizable,

$$A = \begin{bmatrix} -20 & 0 & 1 & 0 & 1 \\ 2 & -10 & 0 & 3 & 0 \\ 0 & 4 & 0 & 4 & 1 \\ 0 & 3 & 0 & 10 & 2 \\ 2 & 0 & 1 & 0 & 20 \end{bmatrix}.$$

3. Let

$$A = \begin{bmatrix} -2 & 1 & 0 \\ 1 & -2 & 1 \\ 0 & 1 & -2 \end{bmatrix}.$$

Approximate the biggest eigenvalue of A using the Power method. Perform three steps of the method with initial vector $x^{(0)} = (1, 1, 1)^T$.

4. Consider the QR-algorithm for a matrix A : Let $A_0 = A$, then

$$A_{k+1} = R_k Q_k, \quad k \in \mathbb{N},$$

where $A_k = Q_k R_k$ is the QR-decomposition of A_k , Q_k is unitary matrix and R_k is right triangular matrix. Show that:

$$(a) A_{k+1} = Q_k^* A_k Q_k$$

$$(b) A_{k+1} = (Q_0 Q_1 \cdots Q_k)^* A (Q_0 Q_1 \cdots Q_k)$$

$$(c) A^k = (Q_0 Q_1 \cdots Q_k) (R_k R_{k-1} \cdots R_0)$$

5. Write a MATLAB-PROGRAM that for a given matrix $A \in \mathbb{C}^{n \times n}$ results in a graph that represents the Gershgorin circles for the matrices A and A^* in the complex plane.
6. Consider a symmetric matrix $A \in \mathbb{R}^{n \times n}$ with n real eigenvalues

$$|\lambda_1| > \dots > |\lambda_n|$$

and corresponding eigenvectors which form an orthonormal basis $\{v_1, \dots, v_n\}$, (with respect to the $\|\cdot\|_2$ norm). Then, the matrix B given by

$$B = A - \lambda_k v_k v_k^T, \quad \text{for a given } k = 1, \dots, n$$

admits the eigenvalues $\lambda_1, \dots, \lambda_{k-1}, 0, \lambda_{k+1}, \dots, \lambda_n$. Implement in MATLAB, a function that approximates all the eigenvalues of A , considering the Power method and the given matrix B .